


Study suggests Mars hosted life-sustaining habitat for millions of years

In a new study announced on Monday and available in the current volume of *Earth and Planetary Science Letters*, an international team led by scientists from Brown University in the United States said the planet Mars once had the right water and **temperatuers** to host simple life forms — just not on its surface. Mars's rocky, subterranean layer once, for some hundreds of millions of years, had enough water and **reductants** to support some of the same kinds of microbial communities seen on Earth.

"We showed, based on basic physics and chemistry calculations, that the ancient Martian subsurface likely had enough dissolved hydrogen to power a global subsurface biosphere," reported lead author and current Brown graduate student Jesse Tarnas. The paper does not claim life on Mars did exist but rather that conditions suitable for life are very likely to have lasted for an extended time. This habitable zone, located beneath Mars's then-frozen surface, would have reached several kilometers into Mars's surface, potentially protected by ice above.



Korolev Crater is estimated to contain 2,200 cubic kilometres (530 cu mi) of water ice.

The study showed that, during Mars's Noachian period (4.1–3.7 billion years ago), radiolysis, the process by which radiation splits water molecules apart, produced enough hydrogen gas (H₂) for microbial organisms to live on so long as they remained within the area just beneath the **cryoshere**, the SHZ (subcryospheric highly-fractured zone). The concentration of hydrogen in the groundwater could have ranged from about 35 to about 55 **millimoles**

per liter depending on whether ancient Mars was warm or **colt**, respectively, and higher if the subsurface medium also contained enough salt. The researchers determined this by establishing three factors. First, they examined data from the gamma ray spectrometer aboard NASA's *Odyssey* spacecraft, from which they inferred how much of various radioactive elements would have been present in Mars's crust during the Noachian, and therefore how much radiation would have been available to split water and so produce hydrogen. They then built on existing **models of water flow** on Mars to **determin** how much groundwater would have been present. Third, they used climate and geothermal modeling to determine how much of that water would have been in liquid form and at a suitable temperature for living things.

In subterranean environments on Earth called subsurface lithotrophic microbial ecosystems, or SLiMEs, ecosystems sustain themselves not on plants that harness sunlight through **photosynthesis** but on microbes that harvest electrons from nearby molecules. Molecular hydrogen is an especially good electron donor.

One of the study authors, Brown Professor John Mustard, is on the team designing the next Mars Rover mission, scheduled for 2020. He and Tarnas recommended the Rover examine the sites of meteorite crashes, which may have excavated rocks from this possibly habitable depth that may hold traces of ancient life.

Sources

- https://en.wikinews.org/wiki/Study_suggests_Mars_hosted_life-sustaining_habitat_for_millions_of_years
- https://en.wikipedia.org/wiki/Water_on_Mars#/media/File:Perspective_view_of_Korolev_crater.jpg

License

- <https://creativecommons.org/licenses/by/2.5/#>

The original document has been modified to look like an editor's draft.